

AD-A031 969

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

POINTWISE CONTRACTION CRITERIA FOR THE EXISTENCE OF FIXED POINT--ETC(U)

JUL 76 F H CLARKE

DAA629-75-C-0024

UNCLASSIFIED

MRC-TSR-1658


NL

1 OF 1  
AD  
A031969



END

DATE  
FILMED  
1-77



ADA031969

MRC Technical Summary Report #1658

POINTWISE CONTRACTION CRITERIA  
FOR THE EXISTENCE OF FIXED POINTS

Frank H. Clarke

See 1473

**Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706**

July 1976

(Received June 21, 1976)

**Approved for public release  
Distribution unlimited**

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

DDC  
NOV 12 1976  
RECEIVED  
C

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

POINTWISE CONTRACTION CRITERIA FOR THE EXISTENCE  
OF FIXED POINTS

Frank H. Clarke<sup>\*</sup>

Technical Summary Report #1658

July 1976

ABSTRACT

We show that, in a complete metric space, every selfmap that is a "weak directional contraction" admits a fixed point.

AMS (MOS) Subject Classification: 47H10

Key Words: contraction, metric convexity, fixed point

Work Unit Number 1 (Applied Analysis)

ACCESSION for	
NTIS	
DDC	
UNCLASSIFIED	White Section <input checked="" type="checkbox"/>
JUSTIFICATION	Gold Section <input type="checkbox"/>
BY	
DISTRIBUTION/AVAILABILITY NOTES	
Disl.	AVAIL. OR NOT SPECIAL
A	

<sup>\*</sup> Department of Mathematics, University of British Columbia, Vancouver, B.C., Canada V6T 1W5.



POINTWISE CONTRACTION CRITERIA FOR THE EXISTENCE  
OF FIXED POINTS

Frank H. Clarke<sup>\*</sup>

1. Introduction

Let  $(X, \rho)$  be a complete metric space, and let a function  $T : X \rightarrow X$  be given. The celebrated contraction principle of Banach asserts that if there exists a number  $\sigma$  in  $(0, 1)$  such that

$$(*) \quad \rho(Tx, Ty) \leq \sigma \rho(x, y) \quad \forall x, y \in X,$$

( $T$  is then said to be a contraction) then  $T$  has a (unique) fixed point; i.e. a point  $x$  such that  $Tx = x$ .

Our purpose is to investigate what can be said if  $(*)$  holds only in some local sense. For example, suppose for each  $x$  in  $X$  there is some neighborhood  $N(x)$  of  $x$  such that

$$(**) \quad \rho(Tx, Ty) \leq \sigma \rho(x, y) \quad \forall y \in N(x).$$

Must  $T$  have a fixed point? That the answer is negative follows from the fact that any function  $T$  satisfies this condition when  $\rho$  is the discrete metric (i.e. when the range of  $\rho$  is  $\{0, 1\}$ ). Thus any such "pointwise" criterion must be accompanied in some way by at least an indirect hypothesis concerning the metric structure.

---

<sup>\*</sup> Department of Mathematics, University of British Columbia, Vancouver, B.C., Canada V6T 1W5.

In the next section we discuss the main result of this paper, a fixed point theorem for "weak directional contractions". The proof of this result is given in § 4, while § 3 is devoted to refinements of the theorem and some related matters.

## 2. Weak directional contractions

Let  $x$  and  $y$  be points in  $X$ . The open interval between  $x$  and  $y$ , denoted  $(x, y)$ , is given by

$$(x, y) = \{z \in X : z \neq x, z \neq y, \rho(x, z) + \rho(z, y) = \rho(x, y)\}.$$

Let  $T : X \rightarrow X$  be a given mapping. We define  $\underline{DT}(x; y)$ , the lower derivate of  $T$  at  $x$  in the direction of  $y$ , as follows:

$$\underline{DT}(x; y) = 0 \text{ if } y = x, \text{ and otherwise}$$

$$\underline{DT}(x; y) = \liminf_{\substack{z \rightarrow x \\ z \in (x, y)}} \rho(Tz, Tx) / \rho(z, x).$$

This has the usual meaning: for each  $\varepsilon > 0$ , we take the infimum of  $\rho(Tz, Tx) / \rho(z, x)$  over those  $z$  in  $(x, y)$  such that  $\rho(x, z) < \varepsilon$  (this is  $+\infty$  if no such  $z$  exist). The limit of these infima is  $\underline{DT}(x; y)$ .

Definition 1.  $T$  is said to be a weak directional contraction if  $T$  is continuous and if there exists a number  $\sigma$  in  $[0, 1)$  such that  $\underline{DT}(x; Tx) \leq \sigma$  for all  $x$  in  $X$ .

Remark 1. Note that in order for  $T$  to be a weak directional contraction, it is necessary that  $(x, Tx)$  contain points arbitrarily near  $x$  whenever  $x \neq Tx$ . Thus if  $\rho$  is the discrete metric, the only weak directional contraction on  $X$  is the identity mapping. This example shows that the fixed point whose existence is asserted in the following theorem need not be unique.

Theorem 1. Every weak directional contraction on a complete metric space has a fixed point.



Remark 2. M. Edelstein [1] [2] has investigated the question of fixed points for mappings which are contractions in a certain local and uniform sense, by adapting the Picard method of successive approximations (which is ineffective in the context of Theorem 1). Other extensions of the contraction principle are possible when a Banach space structure is present; we refer the reader to Chapter 5 of the monograph by D. R. Smart [4]. The following example lies outside the bounds of the results cited above.

Example. Let  $X = \mathbb{R}^2$ , with the norm given by:

$$\|(x, y)\| = |x| + |y|.$$

If  $\rho((x, y), (x', y')) = \|(x - x', y - y')\|$ , then  $(X, \rho)$  is a complete metric space. It is easy to see that the open interval between any two distinct points  $(x_1, y_1)$  and  $(x_2, y_2)$  consists of the closed solid rectangle having the two given points as diagonally opposite corners, with those two points deleted (this reduces to a line segment in the usual sense if  $x_1$  and  $x_2$  or  $y_1$  and  $y_2$  coincide).

We define  $T : X \rightarrow X$  as follows:

$$T(x, y) = (3x/2 - y/3, x + y/3).$$

It is easily seen that  $T$  is not a contraction (even in a local sense).

However,  $T$  is a weak directional contraction. For let  $T(x, y) \neq (x, y)$ .

Then (setting  $T(x, y) = (a, b)$ ) it follows that  $b \neq y$ , so that the open interval between  $(x, y)$  and  $T(x, y)$  contains points of the form  $(x, z)$  with  $z$  arbitrarily close to  $y$ . But for such points we have:

$$\rho(T(x, z), T(x, y)) / \rho((x, z), (x, y)) = 2/3.$$

Note that the fixed points of  $T$  are all the points of the form  $(x, 3x/2)$ ,  $x \in \mathbb{R}$ .



### 3. Other formulations of the theorem

The following extension of Theorem 1 applies to certain cases in which  $DT(x;Tx)$  is not necessarily bounded away from 1.

Theorem 2. Let  $T$  be a continuous selfmap on a complete metric space  $X$  such that  $DT(x;Tx) < 1$  for all  $x$ . Suppose that every sequence  $\{x_n\}$  in  $X$  such that  $DT(x_n;Tx_n)$  is not bounded away from 1 has a cluster point. Then  $T$  has a fixed point.

Remark 3. The example  $X = [1, \infty)$ ,  $\rho =$  Euclidean metric,  $Tx = x + 1/x$  shows that the cluster point condition cannot be dispensed with. To see that Theorem 2 is indeed more general than Theorem 1, consider a differentiable function  $f : [0,1] \rightarrow [0,1]$  such that  $|f'| < 1$  but  $|f'|$  is not bounded away from 1.

A metric space  $X$  is said to be (metrically) convex if  $(x,y) \neq \emptyset$  for every pair  $(x,y)$  of distinct points. A convex subset of a Banach space has this property.

Definition 2.  $T$  is called a pointwise contraction if for some  $\sigma$  in  $[0,1)$  we have, for all  $x$ ,

$$\limsup_{\substack{y \rightarrow x \\ y \neq x}} \rho(Ty, Tx) / \rho(y, x) \leq \sigma.$$

Corollary 1. Every pointwise contraction on a complete convex metric space has a fixed point.

That this follows from Theorem 1 is a consequence of the following:

(a) every pointwise contraction is continuous and (b) in a complete convex

space,  $(x, y)$  contains points arbitrarily near  $x$  whenever  $x \neq y$ .

These imply that a pointwise contraction on a complete convex space is a weak directional contraction.

When the metric space is convex, Corollary 1 affords a criterion which may be easier to verify than the global contraction condition. It suffices, for example, to prove the following "growth condition": for every  $x$  there is a number  $K(x)$  such that for all  $y$  near  $x$ ,

$$\rho(Ty, Tx) \leq \sigma \rho(y, x) + K(x) \rho(y, x)^2.$$

Question: Is every pointwise contraction on a complete convex metric space a global contraction?

#### 4. Proof of the theorems

It suffices to prove Theorem 2. We now state for convenience the following theorem of Ivar Ekeland [3]:

Theorem. Let  $F : X \rightarrow [0, \infty)$  be a continuous function bounded below, and let  $\varepsilon > 0$  be given. Then there is a point  $u$  such that

$$(i) \quad F(u) < \inf_X F + \varepsilon ,$$

$$(ii) \quad F(x) - F(u) \geq -\varepsilon \rho(x, u) \quad \forall x \in X .$$

Let us define  $F : X \rightarrow [0, \infty)$  as follows:

$$F(x) = \rho(Tx, x) .$$

Since  $T$  is continuous, it follows that  $F$  is continuous. Applying Ekeland's theorem, we deduce the existence, for each positive integer  $K$ , of a point  $u_K$  such that

$$(1) \quad F(u_K) < \inf_X F + 1/K ,$$

$$(2) \quad F(x) + \rho(x, u_K)/K \geq F(u_K) \quad \forall x \in X .$$

If for any  $K$  we have  $F(u_K) = 0$ , then  $u_K$  is a fixed point and we are done. So let us suppose that  $F(u_K)$  is positive for each  $K$ .

Claim:  $\underline{DT}(u_K; Tu_K) \geq 1 - 1/K$ .

Since  $u_K \neq T(u_K)$  there exists a sequence  $\{x_n\}$  in  $(u_K, Tu_K)$  such that  $\rho(u_K, x_n)$  converges to 0 as  $n \rightarrow \infty$ , and

$$(3) \quad \lim_{n \rightarrow \infty} \rho(Tx_n, Tu_K) / \rho(x_n, u_K) = \underline{DT}(u_K; Tu_K) .$$



By definition,

$$(4) \quad \rho(u_K, Tu_K) = \rho(u_K, x_n) + \rho(x_n, Tu_K) .$$

We find (in light of (2)):

$$\begin{aligned} \rho(u_K, Tu_K) &\leq \rho(x_n, Tx_n) + \rho(x_n, u_K)/K \\ &\leq \rho(x_n, Tu_K) + \rho(Tu_K, Tx_n) + \rho(x_n, u_K)/K \\ &\leq \rho(x_n, Tu_K) + \underline{DT}(u_K; Tu_K) \rho(x_n, u_K) + o(\rho(x_n, u_K)) + \rho(x_n, u_K)/K , \end{aligned}$$

where  $o(\rho(x_n, u_K))/\rho(x_n, u_K) \rightarrow 0$  as  $n \rightarrow \infty$ .

Combining this with (4), we arrive at:

$$(5) \quad (1 - 1/K) \rho(x_n, u_K) \leq \underline{DT}(u_K, Tu_K) \rho(x_n, u_K) + o(\rho(x_n, u_K)) .$$

Dividing across by  $\rho(x_n, u_K)$  and letting  $n$  tend to  $\infty$ , we obtain the required inequality.

The hypotheses now imply that the sequence  $\{u_K\}$  has a cluster point  $u$ . In view of (1), we have

$$(6) \quad \rho(x, Tx) \geq \rho(u, Tu) \quad \forall x \in X .$$

If  $u = Tu$  we are done, so let us suppose the contrary and show that

(6) leads to a contradiction. Arguing as we did to obtain (5), we obtain a sequence  $\{x_n\}$  in  $(u, Tu)$  such that  $\rho(x_n, u)$  tends to 0 as  $n \rightarrow \infty$ , and

$$\rho(x_n, u) \leq \underline{DT}(u; Tu) \rho(x_n, u) + o(\rho(x_n, u)) ,$$

where  $o(\rho(x_n, u))/\rho(x_n, u) \rightarrow 0$  as  $n \rightarrow \infty$ . This implies

$$\underline{DT}(u; Tu) \geq 1 ,$$

which contradicts the hypotheses. Q. E. D.

#### REFERENCES

1. M. Edelstein, An extension of Banach's contraction principle,  
Proc. Amer. Math. Soc. 12 (1961) 7-10.
2. M. Edelstein, On fixed and periodic points under contractive mappings,  
J. London Math. Soc. 37 (1962) 74-79.
3. I. Ekeland, On the variational principle, J. Math. Anal. Appl. 47  
(1974) 324-353.
4. D. R. Smart, "Fixed Point Theorems", Cambridge Tracts in Mathematics  
66, Cambridge University Press, London (1974).

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 MRC-TSR-1658	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) 6 POINTWISE CONTRACTION CRITERIA FOR THE EXISTENCE OF FIXED POINTS.		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) 10 Frank H. /Clarke		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) 15 DAAG29-75-C-0024
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 9 Technical summary rept.		12. REPORT DATE 17 Jul 1976
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		13. NUMBER OF PAGES 9
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
18. SUPPLEMENTARY NOTES		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) contraction metric convexity fixed point		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) I+ is shown. We show that, in a complete metric space, every selfmap that is a "weak directional contraction" admits a fixed point.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED 221200  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered) LB